

Some discussions on strong interaction

Ning Wu

CCAST (World Lab), P.O.Box 8730, Beijing 100080, P.R.China

and

Division 1, Institute of High Energy Physics, P.O.Box 918-1, Beijing 100039, P.R.China

February 1, 2008

PACS Numbers: 12.38-t, 12.38Aw, 14.70Dj, 11.15-q,

Keywords: quantum chromodynamics, gluons, gauge symmetry, mass, glueball

[Abstract] Using the gauge field model with massive gauge bosons, we could construct a new model to describe the strong interaction. In this new quantum chromodynamics(QCD) model, we will introduce two sets of gluon fields, one set is massive and another set is massless. Correspondingly, there may exists three sets of glueballs which have the same spin-parity but have different masses. A possible new kind of long-range force field is also discussed in this paper.

Now, it is generally believed that the most elementary constituents of matter are elementary fermions, such as quarks and leptons, and elementary bosons, such as photon, gluon and gauge bosons $W^\pm$ and Z etc.. Quarks are elementary constituents of hadrons[1-2]. In order to solve the problem of Fermi statistics of quarks inside hadrons, the concept of color of quarks is introduced to physics[3-5]. It is known that quarks' dynamics obeys $SU(3)_c$ symmetry. Using Yang-Mills gauge field theory[6], physicists have found a quantum theory to describe strong interaction—quantum chromodynamics (QCD). We know that, in QCD, there exists only one kind of gauge field. Because QCD has strict $SU(3)_c$ symmetry, the mass of gauge field is zero.

Using the gauge field model which has a set of massive gauge bosons as well as a set of massless gauge bosons [7], we could construct another QCD to describe strong

interaction. Because of color confinement, colored gluons can not exist in nature in a state of free particle, they exist inside glueballs or hybrid states. Because there exist two sets of gluons, there may exist three sets of glueballs in mass spectrum. They are (g_1g_1) , (g_1g_2) and (g_2g_2) , which have the same spin-parity but different masses. Although all gluons are in color octet, some gluons are colorless. If colorless gluons are not restricted by color confinement, these colorless gluons may exist in a state of free particle and there may exist a new kind of long-range force field [8,11]. In experiment, these free gluons may mix in γ photons. So, physicists will find that γ photons take part in strong interaction [10]. The existence of these free massless gluons may explain the phenomenon that γ photons take part in strong interaction.

The quark fields is denoted as

$$\psi_a(x) \quad , \quad (a = 1, 2, 3), \quad (1)$$

where a is color index and the flavor index is omitted. Define

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix}. \quad (2)$$

All ψ form the fundamental representative space of $SU(3)_c$. In the present theory, two different gauge fields $A_{1\mu}(x)$ and $A_{2\mu}(x)$ will be introduced. The lagrangian of the model is:

$$\begin{aligned} \mathcal{L} = & -\bar{\psi}[\gamma^\mu(\partial_\mu - igA_{1\mu}) + m]\psi \\ & - \frac{1}{4K} Tr(A_1^{\mu\nu} A_{1\mu\nu}) - \frac{1}{4K} Tr(A_2^{\mu\nu} A_{2\mu\nu}) \\ & - \frac{\mu^2}{2K} Tr[(\cos\alpha A_1^\mu + \sin\alpha A_2^\mu)(\cos\alpha A_{1\mu} + \sin\alpha A_{2\mu})] \end{aligned} \quad (3)$$

where

$$\begin{aligned} A_{1\mu\nu} &= \partial_\mu A_{1\nu} - \partial_\nu A_{1\mu} - ig[A_{1\mu}, A_{1\nu}] \\ A_{2\mu\nu} &= \partial_\mu A_{2\nu} - \partial_\nu A_{2\mu} + igtg\alpha[A_{2\mu}, A_{2\nu}] \end{aligned} \quad (4)$$

The above lagrangian can be proved to have strict $SU(3)_c$ gauge symmetry.

In the above lagrangian, $A_{1\mu}$ and $A_{2\mu}$ are not eigenvectors of mass matrix. So, let's make the following transformations:

$$G_{1\mu} = \cos\alpha A_{1\mu} + \sin\alpha A_{2\mu} \quad (5)$$

$$G_{2\mu} = -\sin\alpha A_{1\mu} + \cos\alpha A_{2\mu} \quad (6)$$

Then, the lagrangian given by eq(3) will change into

$$\begin{aligned} \mathcal{L} = & -\bar{\psi}[\gamma^\mu(\partial_\mu - ig\cos\alpha G_{1\mu} + ig\sin\alpha G_{2\mu}) + m]\psi \\ & - \frac{1}{4} G_{10}^{i\mu\nu} G_{10\mu\nu}^i - \frac{1}{4} G_{20}^{i\mu\nu} G_{20\mu\nu}^i - \frac{\mu^2}{2} G_{1\mu}^{i\mu} G_{1\mu}^i + \mathcal{L}_{gI}, \end{aligned} \quad (7)$$

where

$$G_{m0\mu\nu}^i = \partial_\mu G_{m\nu}^i - \partial_\nu G_{m\mu}^i \quad (m = 1, 2) \quad (8)$$

$$G_{m\mu} = G_{m\mu}^i \lambda_i / 2, \quad (9)$$

and $\mathcal{L}_{gI}$ only contains interaction terms of gauge fields. From the above lagrangian, we see that, gluon fields G_1 are massive fields whose masses are μ , and gluon fields G_2 are massless fields. We use g_1 and g_2 denote gluons correspond to the gluon fields G_1 and G_2 respectively. Because of color confinement, all colored gluons must exist in bound states. Two kinds of gluons may form three kinds of glueballs which are denoted by (g_1g_1) , (g_1g_2) and (g_2g_2) . They have the same spin-parity but different masses. Because gluon g_1 is massive and gluon g_2 is massless, glueball (g_1g_1) is heavier than glueball (g_1g_2) and glueball (g_1g_2) is heavier than glueball (g_2g_2) . If $\text{tg}\alpha$ is not very small or very big, then we will see these three kinds of glueballs in the mass spectrum simultaneously. If $\text{tg}\alpha$ is very small, then only the peak corresponds to glueball (g_1g_1) is obvious, and those peaks correspond to glueballs (g_1g_2) and (g_2g_2) are very weak. In this case, only glueball (g_1g_1) can be seen in the mass spectrum. Similarly, if $\text{tg}\alpha$ is very big, only glueball (g_2g_2) can be seen in the mass spectrum.

From eq(7), we know that the coupling constant between massive gluon fields and quark is $-g\cos\alpha$, and the coupling constant between massless gluon fields and quark is $g\sin\alpha$. Because strong interaction is a short range interaction, we anticipate that the coupling between massive gluon fields and quark is much stronger than the coupling between massless gluon fields and quark. Therefore,

$$\text{tg}\alpha \ll 1. \quad (10)$$

And the strong interaction mainly originate from the interchange of massive gluons. The range of strong force depends on the mass of gluon. Suppose that the range of strong force is about the radius of proton, then the mass of gluon is about 1Gev . If this is true, the masses of g_1^3 and g_1^8 are also about 1Gev . In this case, we could prove that, in a proper approximation, the potential of the strong interaction is linear[11].

Because of color confinement, free quarks can not be directly seen by experiment though the masses of quarks are not very big. There are a lot of explanations of quark confinement at present. A possible explanation is to suppose that vacuum is an anticolor medium [9]. According to this point of view, all colored particles will be confined in bag and all particles detected by experiment are colorless particle. Now, let's discuss the problem on gluon confinement. In eq(9), λ_i are Gell-Mann matrix. Substitute the expression of λ_i into eq(9), we obtain

$$G_\mu = \frac{1}{2} \begin{pmatrix} G_\mu^3 + \frac{1}{\sqrt{3}}G_\mu^8 & G_\mu^1 - iG_\mu^2 & G_\mu^4 - iG_\mu^5 \\ G_\mu^1 + iG_\mu^2 & -G_\mu^3 + \frac{1}{\sqrt{3}}G_\mu^8 & G_\mu^6 - iG_\mu^7 \\ G_\mu^4 + iG_\mu^5 & G_\mu^6 + iG_\mu^7 & -\frac{2}{\sqrt{3}}G_\mu^8 \end{pmatrix}. \quad (11)$$

In the above relation, G_μ represents $G_{1\mu}$ or $G_{2\mu}$. All gluons in the non-diagonal position are colored gluons. The interaction terms between diagonal gluons and quarks are

$$\begin{aligned} & -ig'\bar{\psi}_1\gamma^\mu\psi_1(G_\mu^3 + \frac{1}{\sqrt{3}}G_\mu^8) - ig'\bar{\psi}_2\gamma^\mu\psi_2(-G_\mu^3 + \frac{1}{\sqrt{3}}G_\mu^8) \\ & + \frac{2}{\sqrt{3}}ig'\bar{\psi}_3\gamma^\mu\psi_3G_\mu^8 \end{aligned} \quad (12)$$

where

$$\begin{cases} g' = -g\cos\alpha & \text{if } G_\mu \text{ is } G_{1\mu} \\ g' = g\sin\alpha & \text{if } G_\mu \text{ is } G_{2\mu} \end{cases} \quad (13)$$

Because $\overline{\psi}_m \gamma^\mu \psi_m$ ($m = 1, 2$) are colorless, G_μ^3 and G_μ^8 are colorless gluons. If color confinement originate from the property that vacuum is an anticolor medium, colorless particle, such as G_μ^3 or G_μ^8 , can exist in a state of free particle, and therefore can be detected by experiment.

If gluons g_1^3 and g_1^8 can exist in the state of free particle, some of their properties are similar to those of glueballs, especially their decay modes. They are massive vector particles. They carry no electric charge. Because they couple to quarks and massless gluons, there are hadrons and massless, electric neutral vector particles in their decay products. Those massless, electric neutral vector particles may be regarded as γ photon in the experiment. In other words, g_1^3 and g_1^8 may have "radiative" decay modes. Because g_1^3 and g_1^8 don't couple to leptons, there are almost no leptons in their decay products, and they will not be directly produced in the $e^+ + e^-$ collisions, but they will be produced in the strong decay or in the $p + \overline{p}$ collisions or in the course of any other strong interaction. It is very important in both theory and experiment to search for these two kinds of gluons in the experiment. To verify whether g_1^3 and g_1^8 can exist as free particles will conduce to the understanding of the nature of color confinement.

For gluons g_2^3 and g_2^8 , they are massless, electric neutral vector particle, and can be emitted from or be absorbed by general atoms. These properties are the same as those of photon, so in the experiment, they may be regarded as photon. But they have essential differences. It is known that g_2^3 and g_2^8 take part in strong interaction, and photon does not directly take part in strong interaction. If physicists find that photons take part in strong interaction obviously, that means that there are massless, electric neutral vector particles mixed in photons. In other words, those photons take part in strong interaction might be g_2^3 or g_2^8 gluons. Indeed, physicists have found that photons take part in strong interaction [10], it is important to verify whether there are gluons mixed in those photons. Another important difference between g_2^3 or g_2^8 and photon is that g_2^3 and g_2^8 don't interact with charged leptons e, μ and τ , but photons interact with any charged particles include leptons. Finally, gluons have self interaction, but photons have no self interaction. From theoretical point of view, photon is a particle of no structure, it should not have structure functions. From the present experimental results [10], some gluons might mixed in photons, and the so-called structure functions of photon might be those of gluons.

There may exist a kind of long-range force field transmit by massless gluons g_2^3 and g_2^8 . Correspondingly, there may exists a kind of wave transmit by those gluons. Because all colored particles are confined and all matter exists in nature is colorless, the influences of this long-range force field to the motion of matter is very very small.

Reference:

- [1] M.Gell-Mann, Phys.Lett. **8** (1964) 214
- [2] G.Zweig, CERN preprints, Th. 401 and 412, (1964)
- [3] M.Han, Y.Nambu, Phys.Rev. **B139** (1965) 1006
- [4] H.Fritsch, M.Gell-Mann, Proc.XVI Int. Conf. on High Energy Phys., Vol2, P.135 (Chicago)
- [5] H.Fritsch, M.Gell-Mann, H.Leutwyler, Phys.Lett **B97** (1980)365
- [6] C.N.Yang, R.L.Mills, Phys Rev **96** (1954) 191
- [7] Ning Wu, Gauge Field Model With Massive Gauge Bosons, hep-ph/9802236
- [8] Ning Wu, A New Electroweak Model without Higgs particle, hep-ph/9802237
- [9] T.D.Lee, Particle Physics and Introduction to Field Theory, (Harwood Academic Publishers, Amsterdam, 1983)
- [10] There are many papers discuss the photon interaction experimentally. See for example hep-ex /9711005
- [11] Ning Wu,(in preparation)